arXiv:math/0508054v2 [math.PR] 12 Aug 2005

A necessary condition for the uniqueness of the stationary state of a Markov system

Ivan Werner
Email: ivan_werner@pochta.ru

February 2, 2008

Abstract

We continue the study of Markov systems started in [5]. In this paper, we prove a generalization of Breiman's strong low of large numbers [1] which implies a necessary condition for the uniqueness of the stationary state of a Markov system.

MSC: 60J05, 37D35, 37A50, 37H99, 28A80

Keywords: Markov systems, iterated function systems with place-dependent probabilities, random systems with complete connections, g-measures, Markov chains, equilibrium states, fractals.

An introduction to Markov systems can be found in one of the previous papers by the author, e.g. [5], [6], [7], [8], [9].

Let (K,d) be a complete separable metric space. Let $\mathcal{M} := (K_{i(e)}, w_e, p_e)_{e\in E}$ be a finite Markov systems on K (see figure 1) with an invariant Borel probability measure μ. We assume that $K_{i(e)}$ is open in $K = \bigcup_{e\in E} K_{i(e)}$, $w_e|_{K_{i(e)}}$ is continuous and $p_e|_{K_{i(e)}}$ is continuous and bounded away from zero for all $e \in E$. Obviously, these conditions imply the Feller property of the Markov system. We shall denote by $C_B(K)$ the set of all bounded continuous functions on K and by $P(K)$ the set of all Borel probability measures on K. Let $U : C_B(K) \longrightarrow C_B(K)$ be the Markov operator associated with the Markov system, given by

$$Uf := \sum_{e\in E} p_e f \circ w_e \quad \text{for all } f \in C_B(K),$$

and $U^* : P(K) \longrightarrow P(K)$ be its adjoint operator, given by

$$U^*\nu(f) := \int U(f)\, d\nu \quad \text{for all } f \in C_B(K),\ \nu \in P(K).$$



Fig. 1. A Markov system.

We call a Markov system $\mathcal{M}$ *contractive* if and only if there exists $0 < a < 1$ such that

$$\sum_{e\in E} p_e(x)d(w_e(x), w_e(y)) \leq ad(x,y) \text{ for all } x,y \in K_{i(e)},\ e \in E.$$

Let $\Sigma^+ := \{(\sigma_1, \sigma_2, ...) :\ \sigma_i \in E,\ i \in \mathbb{N}\}$ endowed with the product topology of discreet topologies. For $x \in K$, let P_x be the Borel probability measure on Σ^+ given by

$$P_x({}_1[e_1, ..., e_n]) := p_{e_1}(x)p_{e_2} \circ w_{e_1}(x)...p_{e_n} \circ w_{e_{n-1}} \circ ... \circ w_{e_1}(x),$$

for every cylinder set ${}_1[e_1, ..., e_n] := \{\sigma \in \Sigma^+ :\ \sigma_1 = e_1, ..., \sigma_n = e_n\}$, which represents the Markov process generated by the Markov system with the Dirac initial distribution δ_x.

It was shown in [8] that an irreducible contractive Markov system $\mathcal{M}$ with uniformly continuous probabilities $p_e|_{K_{i(e)}}$ has a unique invariant Borel probability measure if $P_x \ll P_y$ for all $x, y \in K_{i(e)}$, $e \in E$, (this was shown in [8] for some locally compact spaces, but it holds also on complete separable spaces, as contractive $\mathcal{M}$ also posses invariant measures on such spaces [2]).

This paper is motivated by the following question. Suppose the Markov system $\mathcal{M}$ has a unique invariant Borel probability measure. Does this imply some restrictions on the measures P_x, $x \in K$? An answer to this question gives a generalization of Breiman's strong low of large numbers [1].

Theorem 1 *Suppose* $1/n\sum_{k=0}^{n-1} U^k g(x) \to \mu(g)$ *for all* $x \in K$, $g \in C_B(K)$. *Let* $f_e : K \longrightarrow [-\infty, +\infty]$ *be Borel measurable such that* $f_e|_{K_{i(e)}}$ *is bounded and* *continuous for all $e \in E$. Then, for every $x \in K$,*

$$\lim_{n\to\infty} \frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ ... \circ w_{\sigma_1}(x) = \sum_{e\in E} \int_{K_{i(e)}} p_e f_e \; d\mu$$

for P_x-a.e. $\sigma \in \Sigma^+$.

Proof. We climb on the shoulders of Breiman [1]. By Kakutani-Yosida norms ergodic lemma (e.g. see [3] p. 441), $1/n \sum_{k=0}^{n-1} U^k g$ converges uniformly to $\int g \; d\mu$ for all $g \in C_B(K)$. Therefore, $1/n \sum_{k=0}^{n-1} U^k (p_e f_e)$ converges uniformly to $\int p_e f_e \; d\mu$ for all $e \in E$ (we use the convention $0 \times \infty = 0$).

Now, fix $x \in K$ and define

$$X_n(\sigma) := \begin{cases} w_{\sigma_n} \circ w_{\sigma_{n-1}} \circ ... \circ w_{\sigma_1}(x), & n \geq 1 \\ x, & n = 0, \end{cases}$$

$$X'_n(\sigma) := f_{\sigma_n} \circ X_{n-1}(\sigma)$$

for all $\sigma \in \Sigma^+$, $n \in \mathbb{N}$,

$$Z_n^1 := \begin{cases} X'_n - E\left(X'_n | X_{n-1}\right), & n > 1 \\ 0, & n \leq 1, \end{cases}$$

and

$$Z_n^k := \begin{cases} E\left(X'_n | X_{n-k+1}\right) - E\left(X'_n | X_{n-k}\right), & n > k \\ 0, & n \leq k \end{cases}$$

for $k > 1$, where E(.|.) denotes the conditional expectation with respect to measure P_x.

We are going to use the following result ([3], p. 387). Let $Y_1, Y_2, ...$ be a sequence of random variables such that $E(Y_n | Y_{n-1}, ..., Y_1) = 0$ and $EY_n^2 \leq c < \infty$ for all n. Then $1/m \sum_{l=1}^m Y_l \to 0$ a.s..

To apply this, note that

$$E(Z_n^k | Z_{n-1}^k, ..., Z_1^k) = E(E(Z_n^k | X_{n-k}, X_{n-k-1}, ..., X_1) | Z_{n-1}^k, ..., Z_1^k),$$

and that, since the $X_1, X_2, ...$ form a Markov chain,

$$E(Z_n^k | X_{n-k}, X_{n-k-1}, ..., X_1) = E(Z_n^k | X_{n-k}) = 0.$$

Furthermore, $E(Z_n^k)^2 \leq 2 \max_{e \in E} ||f_e|_{K_{i(e)}}||_\infty^2$. Thus,

$$\frac{1}{m} \sum_{l=1}^m Z_l^k \to 0 \quad P_x\text{-a.e..} \tag{1}$$

Now, write

$$X'_n - E(X'_n | X_{n-k}) = Z_n^1 + Z_n^2 + ... + Z_n^k, \quad n > k.$$

Thus, by (1),

$$\lim_{m\to\infty}\left|\frac{1}{m}\sum_{n=1}^{m}X'_n-\frac{1}{m}\sum_{n=k+1}^{m}E(X'_n|X_{n-k})\right|=0\quad P_x\text{-a.e..}$$

Or, neglecting at most k terms,

$$\lim_{m\to\infty}\left|\frac{1}{m}\sum_{n=1}^{m}X'_n-\frac{1}{m}\sum_{n=1}^{m}E(X'_{n+k}|X_n)\right|=0\quad P_x\text{-a.e..}$$

So that, for fixed s,

$$\lim_{m\to\infty}\left|\frac{1}{m}\sum_{n=1}^{m}X'_n-\frac{1}{m}\sum_{n=1}^{m}\left(\frac{1}{s}\sum_{k=1}^{s}E(X'_{n+k}|X_n)\right)\right|=0\quad P_x\text{-a.e..}\tag{2}$$

Now, observe that

$$\begin{aligned}&E(X'_{n+k}|X_n)\\ =&\sum_{e_{n+1},\dots,e_{n+k}}p_{e_{n+1}}(X_n)p_{e_{n+2}}(X_n)\dots\\ &\times p_{e_{n+k}}\circ w_{e_{n+k-1}}\circ\dots\circ w_{e_{n+1}}(X_n)f_{e_{n+k}}\circ w_{e_{n+k-1}}\circ\dots\circ w_{e_{n+1}}(X_n)\\ =&\sum_{e_{n+k}}U^{k-1}(p_{e_{n+k}}f_{e_{n+k}})(X_n).\end{aligned}$$

Since $1/s\sum_{k=1}^{s}U^{k-1}(p_ef_e)$ converges uniformly to $\int p_ef_e\ d\mu$ for all $e\in E$, for every $\epsilon>0$, we can choose $s>0$ such that

$$\|\frac{1}{s}\sum_{k=1}^{s}\sum_{e\in E}U^{k-1}(p_ef_e)-\sum_{e\in E}\int p_ef_e\ d\mu\|_\infty<\epsilon.$$

By (2), for such s,

$$\lim_{m\to\infty}\left|\frac{1}{m}\sum_{k=1}^{m}X'_n-\sum_{e\in E}\int p_ef_e\ d\mu\right|\le\epsilon\quad P_x\text{-a.e..}$$

□

Corollary 1 *Suppose K is a compact metric space and μ is a unique invariant Borel probability measure of the Markov system $\mathcal{M}$. Let $f_e:K\to[-\infty,+\infty]$ be Borel measurable such that $f_e|_{K_{i(e)}}$ is continuous for all $e\in E$. Then, for every $x\in K$,*

$$\lim_{n\to\infty}\frac{1}{n}\sum_{k=0}^{n-1}f_{\sigma_{k+1}}\circ w_{\sigma_k}\circ\dots\circ w_{\sigma_1}(x)=\sum_{e\in E}\int_{K_{i(e)}}p_ef_e\ d\mu$$

for P_x-a.e. $\sigma\in\Sigma^+$.

Proof. By the hypothesis $P(K)$ is weakly* compact. Let $x \in K$. Since every weakly* convergent subsequence $1/n_m \sum_{k=0}^{n_m-1} U^{*k}\delta_x$ converges to an invariant Borel probability measure of U^*, by the uniqueness of μ, we conclude that

$$\frac{1}{n}\sum_{k=0}^{n-1} U^{*k}\delta_x \overset{w^*}{\to} \mu.$$

Hence, the claim follows by Theorem 1. □

Remark 1 Obviously, Corollary 1 is a generalization of Breiman's strong low of large numbers [1] for finite Markov systems.

The following is a version for contractive Markov systems.

Corollary 2 *Suppose K is a complete separable metric space. Suppose $\mathcal{M}$ is a contractive Markov system with a unique invariant Borel probability measure μ. Let $f_e : K \to [-\infty, +\infty]$ be Borel measurable such that $f_e|_{K_{i(e)}}$ is bounded and continuous for all $e \in E$. Then, for every $x \in K$,*

$$\lim_{n\to\infty}\frac{1}{n}\sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x) = \sum_{e\in E}\int_{K_{i(e)}} p_e f_e \; d\mu$$

for P_x-a.e. $\sigma \in \Sigma^+$.

Proof. It was shown in [2] that the hypothesis implies

$$\frac{1}{n}\sum_{k=0}^{n-1} U^k g(x) \to \mu(g)$$

for all $x \in K$, $g \in C_B(K)$. Thus, the claim follows by Theorem 1. □

Definition 1 Let $\Sigma := \{(\dots, \sigma_{-1}, \sigma_0, \sigma_1, \dots) : e_i \in E \ \forall i \in \mathbb{Z}\}$ endowed with the product topology of discreet topologies. Denote by S the left shift map on Σ. We call the shift invariant Borel probability measure M on Σ given by

$$M\left({}_m[e_1, \dots, e_k]\right) := \int p_{e_1}(x) p_{e_2}(w_{e_1}x) \dots p_{e_k}(w_{e_{k-1}} \circ \dots \circ w_{e_1}x) d\mu(x),$$

for every cylinder set ${}_m[e_1, \dots, e_k] := \{\sigma \in \Sigma : \ \sigma_i = e_i \text{ for all } m \le i \le n\}$, the *generalized Markov measure* (see [6]). We call the measure preserving transformation $S : (\Sigma, M) \longrightarrow (\Sigma, M)$ a *generalized Markov shift.*

The following corollary gives a necessary condition on measures P_x, $x \in K$, for the uniqueness of μ.

Corollary 3 *Suppose μ is a unique invariant Borel probability measure of $\mathcal{M}$. Suppose K is a compact metric space, or K is a complete separable metric space and $\mathcal{M}$ is contractive. Then, for every $x \in K$,*

$$\lim_{n\to\infty}\frac{1}{n}\log P_x({}_1[\sigma_1,...,\sigma_n]) = \sum_{e\in E}\int_{K_{i(e)}} p_e \log p_e \; d\mu$$

for P_x-a.e. $\sigma \in \Sigma^+$.

Proof. Set $f_e := \log p_e$ for all $e \in E$. Then, by Corollary 1 or Corollary 2, for every $x \in K$,

$$\lim_{n\to\infty}\frac{1}{n}\log[p_{\sigma_1}(x)p_{\sigma_2}\circ w_{\sigma_1}(x)...p_{\sigma_n}\circ w_{\sigma_{n-1}}\circ ...\circ w_{\sigma_1}(x)] = \sum_{e\in E}\int_{K_{i(e)}} p_e \log p_e \; d\mu$$

for P_x-a.e. $\sigma \in \Sigma^+$. $\square$

Remark 2 Note that $-\sum_{e\in E}\int_{K_{i(e)}} p_e \log p_e \; d\mu$ is the Kolmogorov-Sinai entropy of the generalized Markov shift associated with $\mathcal{M}$ and μ (this was proved in [9] under additional assumption of uniform continuity of each $p_e|_{K_{i(e)}}$). This entropy formula also plays a central role in [4].

References

[1] L. Breiman, The strong law of large numbers for a class of Markov chains, *Ann. Math. Statist.* **31** (1960) 801–803.

[2] K. Horbacz and T. Szarek, Irreducible Markov systems on polish spaces, *preprint.*

[3] M. Loève, *Probability Theory*, D. Van Nostrand, New York, 1955.

[4] W. Slomczynski, *Dynamical entropy, Markov operators, and itereated function systems*, Rozprawy Habilitacyjne Uniwersytetu Jagiellońskiego Nr 362, Wydawnictwo Uniwersytetu Jagiellońskiego (2003).

[5] I. Werner, Contractive Markov systems, *J. London Math. Soc.* **71** (2005) 236-258.

[6] I. Werner, Coding map for a contractive Markov system, *Math. Proc. Camb. Phil. Soc.* to appear **140** (2) (2006), *preprint math.PR/0504247.*

[7] I. Werner, The generalized Markov measure as an equilibrium state, *Nonlinearity* **18** (2005) 2261-2274.

[8] I. Werner, Contractive Markov systems II, *preprint math.PR/0503633.*

[9] I. Werner, On coding with Feller contractive Markov systems, *preprint math.DS/0506476.*